

CSIRO Grapevine Rootstocks

© CSIRO Plant Industry April 2014
www.csiro.au



Three rootstocks from the CSIRO grapevine rootstock breeding program are available and have been granted Plant Breeders Rights (PBR). The rootstocks are Merbein 5489, Merbein 5512 and Merbein 6262. Here, we summarise the key characteristics of each of the rootstocks and present a comparison of harvest data, pruning weight and wine data compared to Ramsey and 1103 Paulsen, all with Shiraz in Sunraysia over 3 seasons (mean values for the period 2001-2003).

Merbein 5489

Merbein 5489 is a medium vigour rootstock. Pruning-wood weights from Shiraz grafted to Merbein 5489 were about half of those grafted to Ramsey and 1103 Paulsen in a Sunraysia trial. Despite having medium vigour, the Shiraz vines on Merbein 5489 produced similar yields per vine to those grafted to 1103 Paulsen rootstock.

In screening trials, Merbein 5489 was tolerant of the G1, G4 and G30 strains of phylloxera.

Shiraz vines grafted to Merbein 5489 established in a re-plant soil infested with nematodes showed no ill effects over a period of 20 years. Glasshouse studies have shown a low to moderate level of root-knot nematode reproduction on roots by three pathotypes of *Meloidogyne javanica* and one of *M. incognita*. Long term response at sites with other pathotypes has not been examined.

Based on levels accumulated in leaf petioles at harvest under irrigation with Murray water in Sunraysia, Merbein 5489 had an ability equivalent to 1103 Paulsen for sodium and chloride exclusion.

Shiraz vines grafted on Merbein 5489 had a higher crop water use index (CWUI) than those grafted on 1103 Paulsen in a Sunraysia-based trial. Higher CWUI (crop produced per unit of water used) is an indicator of water use efficiency.

Shiraz on Merbein 5489 (Sunraysia trial) accumulated significantly less potassium in berries leading to a lower juice and wine pH and thus less need for acid adjustments in winemaking.

Wine made using fruit from Shiraz vines grafted to Merbein 5489 in the warm Sunraysia region had higher colour density, lower colour hue (brighter wine) and higher total phenolics than equivalent wines made using fruit of Shiraz on 1103 Paulsen and Ramsey rootstocks

Merbein 5512

Merbein 5512 is a low to medium vigour rootstock. Pruning-wood weights from Shiraz vines grafted to Merbein 5512 were about one-third of those grafted to Ramsey and 1103 Paulsen in a Sunraysia trial. The Shiraz vines on Merbein 5512, with their reduced vigour, produced yields per vine which were 25 and 47% lower than those grafted to 1103 Paulsen and Ramsey rootstocks, respectively.

In screening trials, Merbein 5512 was tolerant of the G4 and G30 strains of phylloxera.

Shiraz vines grafted to Merbein 5512 established in a re-plant soil infested with nematodes showed no ill effects over a period of 20 years. Glasshouse studies have shown a moderate to high level of root-knot nematode reproduction on roots by four pathotypes of *Meloidogyne javanica* and one of *M. incognita*, indicating higher sensitivity than 5489 and 6262. Long term response at sites with other pathotypes has not been examined.

Based on levels accumulated in leaf petioles at harvest under irrigation with Murray water in Sunraysia, Merbein 5512 had an ability equivalent to Ramsey and 1103 Paulsen for chloride and sodium exclusion, respectively.

Shiraz on Merbein 5512 (Sunraysia trial) accumulated significantly less potassium in berries leading to lower juice and wine pH and thus less need for acid adjustments in winemaking.

Wine made using fruit from Shiraz vines grafted to Merbein 5512 in the warm Sunraysia region had higher ionized anthocyanins (the coloured form of anthocyanins, data not shown) and lower colour hue (brighter wine) than equivalent wines made using fruit of Shiraz on 1103 Paulsen and Ramsey rootstocks. Wine from Shiraz on Merbein 5512 also had higher colour density and higher total phenolics than that from Shiraz grafted on Ramsey.

Merbein 6262

Merbein 6262 is a low vigour rootstock. Pruning-wood weights from Shiraz grafted to Merbein 6262 were about one-third of those grafted to Ramsey and 1103 Paulsen in a Sunraysia trial. The Shiraz vines on Merbein 6262, with their reduced vigour, produced yields per vine which were 16 and 41% lower than those grafted to 1103 Paulsen and Ramsey rootstocks, respectively.

In screening trials, Merbein 6262 showed moderate tolerance towards the G4 and G30 strains of phylloxera, indicating a risk of phylloxera developing on the roots, but at relatively low levels compared to *Vitis vinifera*.

Shiraz vines grafted to Merbein 6262 established in a re-plant soil infested with nematodes showed no ill effects over a period of 20 years. Glasshouse studies have shown a nil to low level of root-knot nematode reproduction on roots by four pathotypes of *Meloidogyne javanica* and one of *M. incognita*. Long term response at sites with other pathotypes has not been examined.

Based on levels accumulated in leaf petioles at harvest under irrigation with Murray water in Sunraysia, Merbein 6262 was equivalent to 1103 Paulsen and Ramsey for sodium exclusion. However, it accumulates higher chloride concentrations, e.g. 0.95% dry weight in Shiraz petioles versus 0.34% on Ramsey, and is not recommended for saline situations.

Shiraz on Merbein 6262 (Sunraysia trial) accumulated significantly less potassium in berries leading to lower juice and wine pH and thus less need for acid adjustments in winemaking.

Wine made using fruit from Shiraz vines grafted to Merbein 6262 in the Sunraysia region had higher colour density, lower colour hue (brighter wine) and higher total phenolics than equivalent wines from Shiraz on 1103 Paulsen and Ramsey rootstocks.

Harvest data, pruning and wine data, Sunraysia (2001-03)

Shiraz grafted on rootstock	VIGOUR	HARVEST			WINE				
	Pruning Wt. (kg)	Juice °Brix	Juice pH	Yield (kg)	Acid Added (g/L)	Wine pH	Colour Density (au)	Colour Hue	Total Phenolics (au)
1103 Paulsen	4.2 _c	25.6 _d	4.18 _d	11.8 _b	4.72 _d	3.57 _b	6.26 _{ab}	0.61 _b	53.8 _b
Ramsey	4.3 _c	25.2 _c	4.25 _e	16.7 _c	5.95 _e	3.58 _b	5.84 _a	0.59 _b	49.5 _a
Merbein 5489	2.0 _b	24.3 _b	3.83 _a	13.3 _b	1.90 _a	3.49 _a	7.56 _c	0.54 _a	60.9 _c
Merbein 5512	1.5 _{ab}	24.5 _b	3.91 _b	8.8 _a	2.72 _b	3.48 _a	6.55 _b	0.55 _a	56.8 _b
Merbein 6262	1.2 _a	23.7 _a	3.94 _b	9.9 _a	2.60 _{ab}	3.51 _a	8.17 _d	0.53 _a	62.4 _c

Values with different subscripts are significantly different ($p=0.05$)

Suitability of the rootstocks in various locations and regions with various scion varieties and their tolerance of different pests and diseases and strains thereof is yet to be fully tested. Given that the rootstocks with Shiraz as scion vary in vigour and yield to pruning weight ratio, different management strategies will need to be considered in order to obtain best performance in specific situations.

FOR FURTHER INFORMATION

Peter Clingeffer
e peter.clingeffer@csiro.au
p 08 8303 8721
w www.csiro.au/plantindustry

Disclaimer

CSIRO gives no warranty and makes no representation that the information contained in this document is suitable for any purpose or is free from error. CSIRO and its officers, employees and agents accept no responsibility for any person acting or relying upon the information contained in this document, and disclaim all liability for any loss, damage, cost or expense incurred by reason of any person using or relying on the information contained in this document or by reason of any error, omission, defect, or mis-statement contained therein.

ACKNOWLEDGEMENTS

Phylloxera tolerance testing done with input from DPI, Rutherglen, Victoria



FUNDING CONTRIBUTIONS

